


GB_IE-EN



Talon[®]Soft

Rodenticide

TALON SOFT

Active ingredient: Brodifacoum 0.005% w/w (0.05g/kg)

RODENTICIDE FOR PROFESSIONAL USE AGAINST RATS AND MICE. FOR USE INDOORS.

This product is approved under The Control of Pesticides Regulations 1986 (as amended) for use as directed.

PCS 96894

STATUTORY CONDITIONS OF APPROVAL RELATING TO USE.

FOR USE ONLY BY PROFESSIONAL OPERATORS. FOR INDOOR USE ONLY. FOR USE ONLY IN RODENT CONTROL BY PROFESSIONAL OPERATORS. APPLY AS NECESSARY.

Users must comply in full with all their legal responsibilities under the Control of Pesticides Regulations 1986 (as amended), including complying with all Conditions of Approval.

HSE 9468

READ ALL PRECAUTIONS BEFORE USE.

PRECAUTIONS: To avoid risks to man and the environment, comply with the instructions for use. *The COSHH (Control of Substances Hazardous to Health) Regulations 2002 (as amended) apply to the use of this product at work. *Guidance on the safe use of rodenticides is provided in leaflets AIS 31 and MISC 515 at <http://www.hse.gov.uk/pubns/biopestindex.htm>.

*PREVENT ACCESS TO BAIT by children, birds and non-target animals [particularly dogs, cats, pigs, poultry]. *Keep away from food, drink and animal feedingstuffs. *Avoid excessive contamination of coveralls. *Search for and remove rodent bodies at frequent** intervals during treatment (unless used in sewers). Collect and dispose of the remains of bait and any remaining rodent bodies after treatment (unless used in sewers). *You must ensure that you comply with legislation regarding the correct disposal of waste. For further guidance, contact the Environment Agency or your local Scottish Environment Protection Agency (SEPA) office. (** Intervals will vary depending on formulation and treatment regime; therefore the user should refer to and comply with the manufacturer's guidelines). *HARMFUL TO WILDLIFE. *Wild mammals and birds may be at greater risk if this product is not used in accordance with the label. *In case of accident or if you feel unwell seek medical advice immediately (show the label where possible). *WASH HANDS ANDEXPOSED SKIN before meals and after use. *Keep only in the original container. *Keep locked up.

DISPOSAL: All pesticides must be disposed of as hazardous waste. This product must be disposed of as hazardous waste. Empty containers may be disposed of as controlled waste in accordance with the appropriate regulations. Collect and handle bodies with caution to avoid disease transfer (wear waterproof gloves). Dispose of dead rodents in accordance with local requirements. Remove all baits and bait trays after treatment and dispose of them in accordance with local requirements.

MARKETED BY: Syngenta Crop Protection AG, Schwarzwaldallee 215, Basel, Switzerland Emergency Line: +44 (0)1484 538444

NOTE TO DOCTOR: *Brodifacoum is an indirect anticoagulant. Phytomenadione, Vitamin K1, is antidotal. Determine prothrombin times not less than eighteen hours after consumption. If elevated, administer Vitamin K1 and continue until prothrombin time normalises. Continue determination of prothrombin times for two weeks after withdrawal of the antidote and resume treatment if elevation occurs in that time. For further information (UK) contact the National Poisons Information Service on 0844 892 011 (for use by medical professionals only). For further information (Ireland) contact the National Poisons Information Centre (NPIC): 01 837 9964 or 01 809 2566 (for Health care professionals only, 24 hours). 01 809 2166 (Public poisons information line, 9am-5pm, Monday-Friday).

FIRST AID: This product contains an anticoagulant compound. If ingested, symptoms may include nosebleed and bleeding gums. In severe cases, there may be bruising, haematomas and blood present in the faeces and urine. In case of accident or if you feel unwell, seek medical advice immediately and show the label where possible.

DIRECTIONS FOR USE: Wear gloves and any other suitable personal protective equipment. Cut tapered nozzle to give 1cm opening. Cut off end of tube dome and fit nozzle before applying pressure. When not in use, release pressure to prevent leakage and cover the nozzle using the cap provided. When cutting off the dome end of the tube, take care not to damage the screw thread. Apply bait (up to 50g per rat bait box and up to 8g per mouse bait box) using both corners of the bait box. After application, deposit any 'run-on' by drawing the end of the nozzle against the bait container. Cover the nozzle when not in use and release the caulking gun pressure to prevent leakage. Bait boxes should be placed in protected situations along runs and in areas where rodents are moving, but not near hot pipes, vents etc. Inspect bait boxes frequently and replace until no more is taken. Do not use in open trays. As a guide, place boxes 5 metres apart for high infestations and 10 metres apart for low infestations. Always protect baits well so as to prevent access by children, birds and non-target animals. Inspect baiting points at appropriate intervals and replenish bait as necessary until no more bait is taken. Thereafter, revisit as necessary. For low infestations, baits should be checked within the first 7-10 days, and thereafter revisit as necessary. At heavily infested sites check bait points after 3-4 days and replenish, at the end of the first week a third baiting round may be conducted. Thereafter, check the bait points weekly for curative treatments and every month for preventative treatments and replenish with bait, if necessary. Record bait points and amount of bait set out. If bait take is low but rodent activity remains, rearrange bait points. Throughout the treatment searches for the bodies of dead rodents on site and up to 150m away from the site should be conducted to protect wildlife. The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. When this product is used in domestic premises the householder should be provided with an 'advice to householder' leaflet containing the product name, product HSE number and active substance (Brodifacoum). This leaflet should also contain instructions to prevent access to bait by children and domestic animals, particularly dogs and birds, the information that the product is an anticoagulant and the contact details of the National Poisons Information Service.

Contents: 300 g

CCLT17 Label No.: 2012-110

Product names marked ® or ™ - the ALLIANCE FRAME
the SYNGENTA Logo and the PESTROCCO logo
are Trademarks of a Syngenta Group Company

Black 340 Blue 072

Label Size = 186 x 150 mm

DGC Nil 1