



Talon[®] Waxblocks

Rodenticide

syngenta[®]

Talon Waxblocks

Contains 0.005% w/w Brodifacoum.
Contains the human taste deterrent denatonium benzoate.

STATUTORY CONDITIONS OF APPROVAL RELATING TO USE

FOR USE ONLY IN RODENT CONTROL BY PROFESSIONAL OPERATORS INDOORS

APPLY AS NECESSARY

Users must comply in full with all their legal responsibilities under the Control of Pesticides Regulations 1986 (as amended), including complying with all Conditions of Approval.
READ ALL PRECAUTIONS BEFORE USE.

HSE 9471

Talon Waxblocks contain a potent anticoagulant rodenticide, which is active against rats and mice, including those resistant to some other anticoagulants. A lethal dose can be ingested as only part of a single day's food intake – multiple feeding is not usually necessary. Symptoms do not normally occur for several days, so bait shyness would not be expected. As multiple feeding is not usually necessary, Talon[®] can be used most economically and safely when small quantities are applied with an interval in between application.

The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used.

*When this product is used in domestic premises the householder should be provided with an "Advice to householders" leaflet containing the product name, product HSE number and the active substance: Brodifacoum. This leaflet should also contain instructions to prevent access to bait by children and domestic animals, particularly dogs and birds, the information that the product is anticoagulant and the contact details of the National Poisons Information Service.

2.5 kg

Product names marked ® or ™, the ALLIANCE FRAME, the SYNGENTA Logo and the PURPOSE (C) W are Trademarks of a Syngenta Group Company

DIRECTIONS FOR USE

Place bait in bait boxes or other suitable containers inside infested buildings. Assess the degree of infestation, look for droppings, holes, gnaw marks and tracks, then position bait in areas where rodents are active. Use the higher application rate where rats and mice are numerous and lay baits at the denser spacing i.e. every 5 metre for rats and every 2 metre for mice. Rats generally feed in sheltered positions in surroundings with which they have become familiar. Lay bait along defined runs, near walls or where rats are feeding. Mice are sporadic feeders and will explore new objects so bait positioning should be changed periodically.

Rodents controlled	Amount of 'Talon' per baiting point	Recommended space between baiting points
Rats	1-3 blocks	5-10 metre
Mice	1 block	2.5 metre (move periodically)

Initial inspections of bait points after 3 or 4 days is recommended. Replace any bait that has been consumed. Repeat approximately 4 days later and then weekly. Replace contaminated or spoiled blocks.

NOTE TO DOCTOR

*Brodifacoum is an indirect anticoagulant. Phytonadione, Vitamin K1, is antidotal. Determine prothrombin times not less than eighteen hours after consumption. If elevated, administer Vitamin K1 until prothrombin time normalises.

*Continue determination of prothrombin time for two weeks after withdrawal of antidote and resume treatment if elevation occurs in that time.

PRECAUTIONS

- To avoid risks to man and the environment, comply with instructions for use.
- * The (COSHH) Control of Substances Hazardous to Health Regulations 2002 (as amended) may apply to the use of this product at work.
- * Guidance on the safe use of rodenticides is provided in leaflets AIS 31 and MISC 515 at <http://www.hse.gov.uk/pubns/biopestindex.htm>
- * PREVENT ACCESS TO BAIT by children, birds and non-target animals particularly dogs, cats, pigs and poultry.
- * Avoid excessive contamination of coveralls.

- * Search for and remove rodent bodies at frequent** intervals during treatment. Collect and dispose of the remains of bait and any remaining rodent bodies after treatment.
- * You must ensure that you comply with legislation regarding the correct disposal of waste. For further guidance, contact the Environment Agency or your local Scottish Environment Protection Agency (SEPA) office. (**Intervals will vary depending on formulation and treatment regime; therefore the user should refer to and comply with the manufacturer's guidelines).
- * HARMFUL TO WILDLIFE
- * FOR INDOOR USE ONLY
- * Wild mammals and birds may be at greater risk if this product is not used in accordance with the label.
- * IF YOU FEEL UNWELL, seek medical advice (show the label where possible).
- * WASH HANDS AND EXPOSED SKIN before meals and after use.
- * Keep only in the original container.
- * KEEP IN A SAFE PLACE.
- * Keep away from food, drink and animal feeding stuffs.
- * Where appropriate, secure the bait block at its placement site to reduce the likelihood of its removal by rodents.

This product is approved under the Control of Pesticides Regulations 1986 for use as directed.

HSE 9471

'Talon' is a Trade Mark of Syngenta.

Marketed by

Syngenta Crop Protection
AG, CH 4002, Basel, Switzerland
Telephone: +44(0)1484 538444

TM